$CCG\ Outcome\ Framework-April\ 2017\ -\ \underline{http://ccgtools.england.nhs.uk/ccgoutcomes/flash/atlas.html}$

Indicator	Value		England	Cluster	England Min	Spine chart	England Ma
CCG Outcomes Indicator Set- domain 1							
.1 Potential years of life lost (PYLL) from causes considered amenable to healthcar	1,778 🍩	3	1,869	2,122	1,055	•1 •	3,204
.1 Potential years of life lost (PVLL) from causes considered amenable to healthcar	2,331	1	2,266	2,742	1,325		3,902
.2 Under 75 mortality rates from cardiovascular disease	54.2	4	64.0	7.5.9	37.1	• •	130.2
.3 Completion of cardiac rehabilitation following an admission for coronary heart dis	46.0 🌘	3	38.0	38.7	0.0	* •	75.4
4 Myocardial infarction, stroke and stage 5 chronic kidney disease in people with d	137,9 •		100.0	99.2	59.8	+	202.7
.5 Mortality within 30 days of hospital admission for stroke	1.23 •	*	1.04	1.09	0.43	1 • •	1.63
6 Under 75 mortality rates from respiratory disease	26.3 🌘	1	29.4	38.4	13.6	•1 •	74.3
7 Under 75 mortality rates from liver disease	22.4 •	1	16.1	21.3	7.9	1 •	40.3
.8 Emergency admissions for alcohol related liver disease	51.5		28.2	42,4	3.5	• •	95.4
9 Under 75 mortality rates from cancer	122 🌘	a ·	120	135	90	▶ •	181
10 One-year survival from all cancers	71.9	1	70.4	69.0	64.7	•	74.5
.11 One-year survival from breast, lung and colorectal cancers	71.8	1	69.3	68.7	62.1	• •	76.2
.12 People with Serious Mental Illness (SMI) who have received the complete list o	36.6 🌘		34.8	39.1	17.5	10.	52.4
14 Maternal smoking at delivery	9.00	4	10.38	14.74	1.80	•1 •	31.00
15 Breast feeding prevalence at 6 - 8 weeks	50.4 🌘	1	47.3	34.5	20.7	10	82.5
.17 Record of stage of cancer at diagnosis	76,5 🥮		75.9	75.6	45.1	•	87.8
.18 Percentage of cancers detected at stage 1 and 2	50.8 🌘	*	50.7	49.3	33.3		61.0
.19 Record of lung cancer stage at decision to treat	89.9 🌘	4	90.1	91.5	41.6	•	100.0
.20 Mortality from breast cancer in females	33.4 🌘	*	34.3	34.5	24.0	• •	46,4
.21 All-cause mortality – 12 months following a first emergency admission to hosp	96.4 🌘	4	100.0	103.3	73.7	• 1 •	117.1
22 Hip fracture: incidence	478 🔵		434	480	38	1 @	623
23 Smoking rates in people with serious mental illness (SMI)	39.7 🌘		40.5	43.8	27.2	(•I	55.0
24 Referrals to cardiac rehabilitation within 5 days of an admission for coronary he	40.70	1	14.10	16.75	0.00	1.	41.90
25 Neonatal mortality and stillbirths	6.76	1	7.11	7.39	1.82	0 +	13.18
26 Low birth weight full-term babies	2,30		2.90	3.03	1.30	• 1•	5.80

▼ CCG Outcomes Indicator Set- domain 2						
2.1 Health-related quality of life for people with long-term conditions	0.75 🌖 🛊	0.74	0.71	0.66	• I •	0.82
2.2 Proportion of people who are feeling supported to manage their condition	66.2 🌖 🕴	64.3	64.7	50.5	I+ ●	73.0
2.3 The percentage of people with Chronic Obstructive Pulmonary Disease (COPD) a	12.0 🌘 👚	18.8	19.1	3.8	• •	68.5
2.4 Percentage of people with diabetes who have received nine care processes	54.9 🌷 👢	52.6	54.7	24.5	10	86.3
2.5 People with diabetes diagnosed less than a year referred to structured education	79.0 🌘 🛊	77.2	78.4	41.7		93.2
2.6 Unplanned hospitalisation for chronic ambulatory care sensitive conditions	1,055 • •	816	1,033	73	l 🧑	1,467
2.7 Unplanned hospitalisation for asthma, diabetes and epilepsy in under 19s	438 • •	312	364	24	•	727
2.8 Complications associated with diabetes	123.1 • •	100.0	118.7	63.8	••	164.3
2.9 Access to community mental health services by people from Black and Minority	2,289 • •	2,201	2,082	658	+ 0	5,283
2.10 Access to psychological therapies services by people from Black and Minority	1,585 • 👚	1,312	1,344	295	•	13,013
2.11a Percentage of referrals to Improving Access to Psychological Therapies (IAP	37.0 • 🛊	43.5	41.9	18.7	• •1	60.5
2.11b Percentage of referrals to Improving Access to Psychological Therapies (IAP	52.9 • 🛊	61.8	61.6	33.6	• •	79.8
2.11c Percentage of referrals to Improving Access to Psychological Therapies (IAP	6.20 🌖 🕴	6.20	6.54	3.40	••	11.30
2.15 Health-related quality of life for carers, aged 18 and above	0.81 🍑 🕴	0.80	0.78	0.70	• I •	0.86
2.16 Health-related quality of life for people with a long-term mental health condition	0.57 • 🛊	0.53	0.49	0.37	+ I •	0.71
▼ CCG Outcomes Indicator Set- domain 3						
3.1 Emergency admissions for acute conditions that should not usually require hospi	1,830 • 👚	1,348	1,725	216	••	2,341
.2 Emergency readmissions within 30 days of discharge from hospital	12.5 🌖 🔻	11.9	12.4	8.9	49	14.5
.3 Elective Hip replacement (Primary) procedures - patient reported outcomes mea	0.45 • 1	0.43	0.43	0.35	+ I •	0.48
.3 Elective knee replacement (Primary) procedures - patient reported outcomes me	0.35 • -	0.31	0.31	0.18	+ •	0.38
3.3 Elective groin hernia procedures - patient reported outcomes measures (PROMS)	0.10 🌑 🛊	0.08	0.08	-0.01	4 •	0.13
.3 Elective varicose veins procedures - patient reported outcomes measures (PRO	n/a	0.10	0.10	0.00	+	0.15
3.4 Emergency admissions for children with lower respiratory tract infections	458 🌖 🐧	457	546	42	• •	936
3.5 People who have had a stroke who are admitted to an acute stroke unit within 4 h	74.4 • 1	60.3	63.1	18.8	1+	85.6
8.6 People who have had an acute stroke who receive thrombolysis	9.10 • 1	11.20	10.22	2.90	0+	30.40
3.7 People with stroke who are discharged from hospital with a joint health and socia	95.1 🌘 👚	87.7	89.5	12.5	№ •	100.0
.8 People who have a follow-up assessment between 4 and 8 months after initial ad	5.00 🌑 👚	29.70	37.93	0.00	• I +	97.30
8.9 People who have had an acute stroke who spend 90% or more of their stay on a	93.9 • 👚	84.1	85.1	61.3	+	95.8
3.10.i Hip fracture: proportion of patients recovering to their previous levels of mobil	n/a	37.4	34.6	0.0	+1	88.9
3.10.ii Hip fracture: proportion of patients recovering to their previous levels of mobi	n/a	61.4	60.8	37.0	4	100.0
.11 Hip fracture: collaborative orthogeriatric care	95.6 🌘 🛊	95.7	95.2	37.9	•	100.0
.12 Hip fracture: timely surgery	80.8 • 1	74.6	72.6	42.7	•1	92.2
.13 Hip fracture: multifactorial falls risk assessment	100.0 🌑 🛕	98.8	98.5	69.1	*	100.0
.14 Alcohol-specific hospital admissions	177.2 • *	114.3	162.9	13.3	••	333.1
8.15 Emergency alcohol-specific readmission to any hospital within 30 days of disc	150.7 • *	100.0	105.8	28.5	1+	199.9
8.16 Unplanned readmissions to mental health services within 30 days of a mental h	134.1 • 1	100.0	94.9	20.9	4	317.5
3.17 Percentage of adults in contact with secondary mental health services in emplo	2.00 • ↓	5.00	4.17	0.00	• •1	18.00
3.18 Hip fracture: care process composite indicator	72,3 🌘 👚	65.6	62.6	11.4	+ •	90.0

